

**Ceratium axiale** Kofoid

*Bull. Mus. Compar. Zool. Harvard Coll.* 50: 170, pl. 4, fig. 26 (1907)

*Neoceratium axiale* (Kofoid) F.Gómez, D.Moreira & López-García, *Protist* 161: 43 (2010), *nom. illeg.*

Type locality: eastern tropical Pacific Ocean

Australian distribution: Qld (Great Barrier Reef), N.S.W. (SE), Tas. (W of King Island)

Extra-Australian distribution: Gulf of Mexico, tropical Atlantic Ocean, tropical Indian Ocean, Solomon Islands, South China Sea, East China Sea, eastern tropical Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)