

Ceratium declinatum G.Karst.

Wiss. Ergeb. Deutschen Tiefsee-Exped. "Valdivia" 2(2): 406, pl. 48, fig. 2 (1907)

Type locality: southern Indian Ocean

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Qld (Great Barrier Reef), N.S.W. (N coast to Sydney), East Australian Current

Extra-Australian distribution: Brazil, Gulf of Mexico, South Atlantic Ocean, North Atlantic Ocean, Canary Islands, British Isles, Bay of Biscay, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, SE Africa, southern Indian Ocean, Sulu Sea, East China Sea, Sea of Japan, North Pacific Ocean, Solomon Islands, Tasman Sea, New Zealand, SW Pacific Ocean, Peru, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)