

Ceratium geniculatum (Lemmerm.) Cleve

Göteborgs Kungl. Vetensk. Samhälle Handl. 4(3): 221 (1901)

Ceratium fusus var. *geniculatum* Lemmerm., *Abh. Naturwiss. Vereine Bremen* 16: 349, pl. 1, fig. 17 (1899)

Type locality: Atlantic Ocean (14°S, 27°W)

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Qld (Great Barrier Reef), N.S.W. (Crowdy Head), East Australian Current

Extra-Australian distribution: Brazil, Gulf of Mexico, U.S.A. (Virginia), Atlantic Ocean, Canary Islands, Madeira, East Africa, Indian Ocean, Bay of Bengal, Indonesia, South China Sea, East China Sea, SW Pacific Ocean, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)