

Ceratium gravidum Gourret

Ann. Mus. Hist. Nat. Marseille (Zool.) 1(8): 58, pl. 1, fig. 15 (1883)

Neoceratium gravidum (Gourret) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (France)

Ceratium gravidum var. *elongatum* E.J.F.Wood, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 40, fig. 146 (1963)

Type locality: Australia (Coral Sea, N.S.W.)

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, N.S.W. (Jibbon Cape to Eden), Vic. (S of Port Fairy), Bass Strait

Extra-Australian distribution: Brazil, Gulf of Mexico, Caribbean Sea, Atlantic Ocean, Canary Islands, British Isles, Spain, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, Indian Ocean, Bay of Bengal, Andaman Sea, Indonesia, Gulf of Thailand, South China Sea, East China Sea, Japan, Pacific Ocean, Solomon Islands, New Zealand

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)