

Ceratium humile Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl.: 40, figs 82, 83 (1911)

Type locality: Japan

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Qld (Great Barrier Reef), East Australian Current, Tas. (Storm Bay)

Extra-Australian distribution: Brazil, Gulf of Mexico, Atlantic Ocean, Red Sea, Arabian Sea, Bay of Bengal, Indonesia, Malaysia, South China Sea, East China Sea, Japan, North Pacific Ocean, Solomon Islands, SW Pacific Ocean, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)