

Ceratium lineatum (Ehrenb.) Cleve

Kongl. Svenska Vetensk. Acad. Handl. 32: 36 (1899)

Peridinium lineatum Ehrenb., *Abh. Königl. Akad. Wiss. Berlin* 1833: pl. 35a, fig. 24c (1834); *Neoceratium lineatum* (Ehrenb.) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: Atlantic Ocean

Australian distribution: N.S.W. (off Sydney), East Australian Current, Bass Strait, Tas. (Storm Bay, off E coast)

Extra-Australian distribution: South Atlantic Ocean, Brazil, Gulf of Mexico, Canary Islands, U.S.A. (NE coast), Canada (E coast), North Atlantic Ocean, Iceland, British Isles, Bay of Biscay, North Sea, Baltic Sea, Norwegian Sea, Barents Sea, Mediterranean Sea, Black Sea, Arabian Sea, Yellow Sea, East China Sea, Sea of Japan, North Pacific Ocean, Bering Sea, Tasman Sea, New Zealand, Southern Ocean, Pacific coast of Mexico, Chile

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

A.McMinn & F.J.Scott, Dinoflagellates, *Antarctic Marine Protists* 202–250 (2005)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)