

Ceratium longirostrum Gourret

Ann. Mus. Hist. Nat. Marseille (Zool.) 1(8): 55, pl. 4, fig. 65 (1883)

Neoceratium longirostrum (Gourret) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (France).

Australian distribution: Qld (Great Barrier Reef), N.S.W. (near Sydney), Bass Strait

Extra-Australian distribution: Brazil, Gulf of Mexico, Caribbean Sea, U.S.A. (Washington), Azores, Canary Islands, West Africa, North Sea, Spain, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, Indian Ocean, Bay of Bengal, South China Sea, East China Sea, Sea of Japan, western Pacific Ocean, North Pacific Ocean, Coral Sea, Tasman Sea, New Zealand, SW Pacific Ocean, eastern tropical Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)