

Ceratium minutum Jörg.

in Schmidt, *Rep. Danish Oceanogr. Exped. Mediterr. Adjacent Seas*, 2, J.1: 24, figs 21–23 (1920)

Type locality: France (Atlantic coast), Spain (Atlantic coast)

Australian distribution: Qld (Great Barrier Reef), N.S.W. (off Sydney), East Australian Current, Tas. (off E coast); “... rather rare from Babel I. [Bass Strait] to Fraser I. [Qld] ... and also on the southern and western coasts of Australia” [Wood, 1954]

Extra-Australian distribution: Gulf of Mexico, Bermuda, U.S.A. (Virginia, Maine), Canada (Bay of Fundy), North Atlantic Ocean, Iceland, Barents Sea, British Isles, North Sea, France (Atlantic coast), Spain (Atlantic coast), Mediterranean Sea, Black Sea, Arabian Sea, North Pacific Ocean, south-western Pacific Ocean, southern Chile

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)