

Gonyaulax digitalis (C.H.G.Pouchet) Kofoid

Univ. Calif. Publ. Zool. 8: 214, pl. 9, figs 1–5 (1911)

Protoperidinium digitale C.H.G.Pouchet, *J. Anat. Physiol.* 19: 443, pl. 18, fig. 14; pl. 19 (1883)

Type locality: France (Atlantic coast)

Australian distribution: Qld (Moreton Bay), N.S.W. (Trial Bay, Port Hacking, Eden), Bass Strait, Tas. (& cysts), W.A. (Swan River)

Extra-Australian distribution: Atlantic Ocean, France, Mediterranean Sea, Pacific Ocean, Southern Ocean (South Georgia, S of Australia), Antarctica (Ross Sea)

Note: Cysts reported from Tas. by McMinn *et al.* (2010) under the palaeontological name, *Bitectatodinium tepikiense* G.J.Wilson

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

A.McMinn & F.J.Scott, Dinoflagellates, *Antarctic Marine Protists* 202–250 (2005)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)