

Gonyaulax polygramma F.Stein

Der Organismus der Infusionstiere 3(2): pl. 4, fig. 15 (1883)

Type locality: Atlantic Ocean, “Südsee”

Australian distribution: Gulf of Carpentaria/North-west Shelf, N.S.W. (coastal and in the East Australian Current), Vic. (Port Fairy), Tas.

Extra-Australian distribution: U.S.A. (Florida), Canary Islands, British Isles, Adriatic Sea, Black Sea, South Africa, South Korea, Southern Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

P.Adjani, T.Ingleton, T.Pritchard & L.Armand, Microalgal blooms in the coastal waters of New South Wales, Australia, *Proc. Linn. Soc. New South Wales* 133: 15–31 (2011)