

Gymnodinium microreticulatum Bolch & Hallegr.

in C.J.S.Bolch, A.Negri & G.M.Hallegraeff, *Phycologia* 38: 303, figs 1–18 (1999)

Type locality: Australia (N.S.W.)

Australian distribution (cysts): Qld (Mackay, Hay Point), N.S.W. (Newcastle), Tas. (SE coast), S.A. (Port Lincoln), W.A. (Port Hedland, Bunbury Harbour, Albany)

Extra-Australian distribution: Uruguay, Portugal, China (Hong Kong), Japan, New Zealand (“cf.”)

References: C.J.S.Bolch, A.P.Negri & G.M.Hallegraeff, *Gymnodinium microreticulatum* sp. nov. (Dinophyceae): a naked, microreticulate cyst-producing dinoflagellate, distinct from *Gymnodinium catenatum* and *Gymnodinium nolleri*, *Phycologia* 38: 301–313 (1999)

C.J.S.Bolch & M.J.Reynolds, Species resolution and global distribution of microreticulate dinoflagellate cysts, *J. Plankton Res.* 24: 565–578 (2002)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)