

**Gymnodinium nanum** J.Schiller

*Arch. Protistenk.* 62: 142, pl. 5, fig. 17 (1928)

*Type locality:* Adriatic Sea

*Australian distribution:* Gulf of Carpentaria/North-west Shelf, Coral Sea, N.S.W. (off Sydney), East Australian Current

*Extra-Australian distribution:* Gulf of Mexico, Spain, Mediterranean Sea (Italy), Adriatic Sea, Black Sea, Indian Ocean, Mexico (Pacific coast)

*References:* E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

S.W.Jeffrey & G.M.Hallegraeff, Studies of phytoplankton species and photosynthetic pigments in a warm core eddy of the East Australian Current. I. Summer populations, *Mar. Ecol. Prog. Ser.* 3: 285–294 (1980)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)