

Gymnodinium venator Flø Jørg. & Sh.Murray

in M.F.Jørgensen, S.Murray & N.Daugbjerg, *J. Phycol.* 40: 1181 (2004)

Type locality: Great Britain

Amphidinium pellucidum Herdman, *Proc. Trans. Liverpool Biol. Soc.* 36: 27, fig. 7 (1922); *Gymnodinium pellucidum* (Herdman) Flø Jørg. & Sh.Murray, in M.F.Jørgensen, S.Murray & N.Daugbjerg, *J. Phycol.* 40: 362 (2004), nom. inval., non *G. pellucidum* Wulff (1919)

Type locality: Great Britain

Amphidinium subsalsum Biecheler, *Bull. Biol. France Belgique* 36 (Suppl.): 6, 27, fig. 6 (1952)

Type locality: France

Australian distribution: N.S.W. (Botany Bay, Narrabeen Lagoon, Port Stephens)

Extra-Australian distribution: Canada (British Columbia), U.S.A. (E coast), British Isles, France, Belgium, Wadden Sea (Denmark), Japan

References: M.F.Jørgensen, S.Murray & N.Daugbjerg, *Amphidinium* revisited: I. Redefinition of *Amphidinium* (Dinophyceae) based on cladistic and molecular phylogenetic analyses, *J. Phycol.* 40: 351–365 (2004)

M.F.Jørgensen, S.Murray & N.Daugbjerg, Corrigendum to: *Amphidinium* revisited: I. Redefinition of *Amphidinium* (Dinophyceae) based on cladistic and molecular phylogenetic analyses, *J. Phycol.* 40: 1181 (2004)

S.Murray, Benthic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 213–259 (2010)