

**Heterodinium milneri** (G.Murray & Whitting) Kofoid

*Univ. Calif. Publ. Zool.* 2: 342 (1906)

*Peridinium milneri* G.Murray & Kofoid, in Murray & Whitting, *Trans. Linn. Soc. London, Ser. Bot.*, 5: 327, pl. 29, fig. 3 ('1899') [1900]

*Type locality:* Atlantic Ocean

*Australian distribution:* Gulf of Carpentaria/North-west Shelf, Coral Sea

*Extra-Australian distribution:* Gulf of Mexico, Canary Islands, Atlantic Ocean, British Isles, Spain, Mediterranean Sea (Turkey; "cf."), Indian Ocean, Pacific coast of Mexico

*References:* E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)