

Mesoporos perforatus (Gran) Lillick

Biol. Bull. Mar. Biol. Lab. Woods Hole 73: 497 (1937)

Exuviaella perforata Gran, *Bull. Plank.* 1912: 99, fig. 7 (1915); *Porella perforata* (Gran) J.Schiller, *Arch. Protistenk.* 53: 55, fig. 12 (1928)

Type locality: North Sea

Australian distribution: N.S.W. (Port Hacking, off Sydney), W.A. (Perth)

Extra-Australian distribution: Gulf of Mexico, Venezuela, U.S.A. (E coast), Canary Islands, British Isles, North Sea, Arctic Ocean, Mediterranean Sea (France, Italy, Greece), Adriatic Sea, Black Sea, Southern Ocean (S of Africa, S of Australia)

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

A.McMinn & F.J.Scott, Dinoflagellates, *Antarctic Marine Protists* 202–250 (2005)