

Ornithocercus heteroporus Kofoid

Univ. Calif. Publ. Zool. 3: 206, fig. 70 (1907)

Type locality: eastern tropical Pacific Ocean

Ornithocercus triclavatus E.J.F.Wood, *Austral. J. Mar. Freshw. Res.* 5: 210 (1954)

Type locality: Australia (N.S.W.)

Ornithocercus biclavatus E.J.F.Wood, *Austral. J. Mar. Freshw. Res.* 5: 211 (1954)

Type locality: Australia (N.S.W.)

Australian distribution: Gulf of Carpentaria/North-west Shelf, N.S.W. (N of Sydney)

Extra-Australian distribution: Brazil, Gulf of Mexico, Canary Islands, Mediterranean Sea, Arabian Sea, Indian Ocean, eastern tropical Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)