

Ornithocercus quadratus F.Schütt

Bot. Zeitung 58(2,16/17): 254, figs 1–4 (1900)

Type locality: Not designated

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld (Great Barrier Reef, Moreton Bay), N.S.W. (E coast), Vic. (W coast), W of Bass Strait

Extra-Australian distribution: Brazil, Caribbean Sea, Gulf of Mexico, Canary Islands, Mediterranean Sea, Arabian Sea, Bay of Bengal, Andaman Sea, South China Sea, East China Sea, Japan

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)