

Oxytoxum milneri G.Murray & Whitting

Trans. Linn. Soc. London, Ser. Bot., 5: 328, pl. 27, fig. 6 ('1899') [1900]

Type locality: Atlantic Ocean

Oxytoxum subulatum Kofoid, *Bull. Mus. Compar. Zool. Harvard Coll.* 50: 190 (1907)

Type locality: eastern tropical Pacific Ocean

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Tasman Sea, N.S.W. (Trial Bay, off Sydney), Tas. (Bass Str.), Indian Ocean (off W.A.)

Extra-Australian distribution: subtropical Atlantic Ocean, Mediterranean Sea, Indian Ocean, eastern tropical Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

J.D.Dodge & R.D.Saunders, A partial revision of the genus *Oxytoxum* (Dinophyceae) with aid of scanning electron microscopy, *Bot. Mar.* 18: 99–122, figs 53–55 (1985)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)