

Phalacroma cuneus F.Schütt

Ergebn. Plankton-Exped. Humboldt-Stiftung 4: 148, pl. 3, fig. 14 (1895)

Dinophysis cuneus (F.Schütt) T.H.Abé, *Publ. Seto Mar. Biol. Lab.* 15(1): 67 (1967)

Type locality: Atlantic Ocean

Phalacroma triangulare E.J.F.Wood, *Austral. J. Mar. Freshw. Res.* 5: 187 (1954); *Dinophysis triangulare* (E.J.F.Wood) Balech, *Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.)* 2: 84 (1967)

Type locality: "East Australia"

Australian distribution: Gulf of Carpentaria, Coral Sea, N.S.W. (as far south as Eden), East Australian Current, North-west Shelf

Extra-Australian distribution: Brazil, Gulf of Mexico, Canary Islands, Atlantic Ocean, Mediterranean Sea, Black Sea, Arabian Sea, Indian Ocean, South China Sea, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)