

Prorocentrum compressum (Bailey) Abé ex J.D.Dodge

J. Linn. Soc., Bot. 71: 110 (1975)

Pyxidicula compressa Bailey, *Smithsonian Contrib. Knowl.* 2(8): 40, pl. 2, figs 13, 14 (1851); *Exuviaella compressa* (Bailey) Ostenf., *Iagttagelser over overfladevants Temperature, Saltoldighet og Plankton paa Islandske og Grönlandske Skibsrouter I* 1899 59 (1899)

Type locality: U.S.A. (Florida)

Australian distribution: Qld (Gulf of Carpentaria; cysts), N.S.W. (Broken Bay, Port Hacking), Vic. (Port Phillip Bay; cysts), Tas. (River Derwent, Tasman Peninsula), W.A. (Fremantle)

Extra-Australian distribution: Brazil, Gulf of Mexico, Caribbean Sea, U.S.A. (Florida, Virginia), Canada (Bay of Fundy), North Atlantic Ocean, Canary Islands, Spain, British Isles, Mediterranean Sea, Adriatic Sea, Black Sea, Arabian Sea, East China Sea, Japan, Pacific Ocean, New Zealand, Southern Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

F.E.Hoedt, J.H.Choat, J.J.Cruz, K.M.Neil & J.Collins, *Survey of the Port of Weipa: Port Marine Baseline Surveys and Surveys for Introduced Marine Pests*. Report by the School of Marine Biology, James Cook University, Townsville (2001)

A.McMinn & F.J.Scott, Dinoflagellates, *Antarctic Marine Protists* 202–250 (2005)

S.Murray, C.L.-C.Ip, R.Moore, Y.Nagahama & Y.Fukuyo, Are prorocentroid dinoflagellates monophyletic? A study of 25 species based on nuclear and mitochondrial genes, *Protists* 160: 245–264 (2009)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)