

**Protoperidinium compressum** (T.H.Abé) Balech

*Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.)* 4: 56 (1974)

*Congruentidium compressum* T.H.Abé, *Sci. Rep. Tohoku Imp. Univ., Biol. Ser.* 4, 2: 420, fig. 36a–e (1927)

Type locality: Japan

Australian distribution: Qld (Gulf of Carpentaria), N.S.W., Vic. (Port Phillip Bay), Tas. (cysts)

Extra-Australian distribution: South Africa, Gulf of Mexico, Baltic Sea, Mediterranean Sea (Italy), Black Sea, India (W coast), South China Sea, Japan

References: C.J.Bolch & G.M.Hallegraeff, Dinoflagellate cysts from Recent marine sediments of Tasmania, Australia, *Bot. Mar.* 33: 173–192 (1990)

J.A.Sonneman & D.R.A.Hill, A taxonomic survey of cyst-producing dino-flagellates from the coastal waters of Victoria, Australia, *Bot. Mar.* 40: 149–177 (1997)

F.E.Hoedt, J.H.Choat, J.J.Cruz, K.M.Neil & J.Collins, *Survey of the Port of Weipa: Port Marine Baseline Surveys and Surveys for Introduced Marine Pests*. Report by the School of Marine Biology, James Cook University, Townsville (2001)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)