

Chaetoceros didymus Ehrenb.

Ber. Bekanntm. Verh. Königl. Preuss. Akad. Wiss. Berlin 1845: 75 (1845)

Type locality: West Africa (?) [in guano]

Australian distribution: N & NW Australia, N.S.W. (North Coast, Sydney), Tas., W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 40 (2010)

Extra-Australian distribution: Argentina, Gulf of Mexico, U.S.A. (Washington, NE coast), Canada (Nova Scotia), British Isles, North Sea, Canary Islands, Spain, Portugal, West Africa, Sea of Japan, North Pacific Ocean, Bering Sea, western, central and southern Pacific Ocean, Southern Ocean