

Chaetoceros peruvianus Brightw.

Quart. J. Microscop. Soc. 4: 107, pl. 7, figs 16–18 (1856)

Type locality: Peru

Australian distribution: N & NW Australia, Qld (Moreton Bay), N.S.W. (North Coast, Central Coast, South Coast), Vic., Tas. (Bass Str.), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 49 (2010)

Extra-Australian distribution: Argentina, Falkland Islands, U.S.A. (E coast), Iceland, British Isles, North Sea, Mediterranean Sea, Black Sea, Canary Islands, South Atlantic Ocean, South Africa, S Indian Ocean, Sea of Japan, Timor Sea, western, south-western, northern and central Pacific Ocean, Pacific coast of South America, Southern Ocean, Antarctica