

Gyrosigma fasciola (Ehrenb.) J.W.Griff. & Henfr.

Diatomaceae 303, pl. 11, fig. 21 (1856)

Ceratoneis fasciola Ehrenb., *Abh. Königl. Akad. Wiss. Berlin* 1839: 147 (1841)

Pleurosigma fasciolum (Ehrenb.) W.Sm., *Ann. Mag. Nat. Hist.*, ser. 2, 9: 9, pl. 2, fig. 6 (1852)

Type locality: Germany (North Sea)

Australian distribution: N.S.W. (Port Hacking), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally epontic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

Extra-Australian distribution: Gulf of Mexico, Caribbean Sea, U.S.A. (Washington, NE coast), Canada (New Brunswick, Nova Scotia), British Isles, North Sea, Baltic Sea, Spain, Adriatic Sea, Romania, Black Sea, Antarctica (King George Island)

Note: In estuarine and marine habitats