

Licmophora ehrenbergii (Kütz.) Grunow f. **ehrenbergii**

Reise Novara, Bot., 1: 36 (1867)

Podosphenia ehrenbergii Kütz., *Kieselschaligen Bacillarien Diatomeen* 121, pl. 9, fig. 13 (1844)

Type locality: North Sea, Baltic Sea, Mediterranean Sea, Adriatic Sea

Australian distribution: Qld (Noosa), Vic. (Gippsland Lakes, Mallacoota Inlet, Port Phillip Bay), Tas. (Georges Bay), W.A. (Perth)

References: N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 100 (2010)

Extra-Australian distribution: California, Pacific coast of Mexico, Caribbean Sea, U.S.A. (Virginia), Canary Islands, British Isles, North Sea, Baltic Sea, White Sea, Barents Sea, Mediterranean Sea (France), Adriatic Sea, Black Sea, Japan