

Plagiotropis lepidoptera (W.Greg.) Kuntze

Revis. Gen. Pl. 3(3): 421 (1898)

Amphiprora lepidoptera W.Greg., *Trans. Roy. Soc. Edinburgh* 21: 505, pl. 12, fig. 59b (1857)

Tropidoneis lepidoptera (W.Greg.) Cleve, *Kongl. Svenska Vetensk.-Akad. Handl.*, ser. 2, 26: 25 (1894)

Type locality: Great Britain

Australian distribution: Qld (Moreton Bay), N.S.W. (Port Stephens, Lake Macquarie, Port Jackson, Botany Bay, Port Hacking, Eden), Vic. (Mallacoota, Gippsland Lakes, Port Phillip Bay), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally eponitic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

Extra-Australian distribution: S Brazil, Gulf of Mexico, Caribbean Sea, U.S.A. (Virginia), Canary Islands, British Isles, North Sea, Baltic Sea, White Sea, Mediterranean Sea (Italy), Adriatic Sea, Romania, Black Sea, Persian Gulf, New Zealand

Note: In estuarine and marine habitats