

Rhizosolenia setigera Brightw.

Quart. J. Microscop. Soc. 6: 95, pl. 5, fig. 7 (1858)

Type locality: Great Britain

Australian distribution: N & NW Australia, Qld (Torres Str., Great Barrier Reef), N.S.W. (North Coast, Central Coast, South Coast), Vic., Tas. (River Derwent), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 31 (2010)

Extra-Australian distribution: Argentina, U.S.A. (Washington, NE coast), Canada (Nova Scotia), British Isles, North Sea, Canary Islands, India, Thailand, Southern Ocean