

Alexandrium catenella (Whedon & Kofoid) Balech

Toxic Dinoflagellates 38, fig. 6 (1985)

Gonyaulax catenella Whedon & Kofoid, *Univ. Calif. Publ. Zool.* 41: 25–31, figs 1–7, 14, 15 (1936)

Type locality: U.S.A. (California)

Australian distribution: N.S.W. (Central & South Coasts), known from Port Hacking (since 1954), Sydney Harbour (1993), Newcastle and Batemans Bay. Also in Port Phillip Bay, Vic. (since 1986), Port River, S.A. (since 1997) and the east coast of Tas. [Hallegraeff *et al.*, 2010]

Extra-Australian distribution: Argentina, Chile, U.S.A. (Alaska, California), British Isles, Mediterranean Sea (Spain, Tunisia, France, Italy), South Africa, China, Japan, South Korea, New Zealand

Note: Produces paralytic shellfish poisons [Hallegraeff *et al.*, 2010]

References: G.M.Hallegraeff, C.J.[S.]Bolch, S.I.Blackburn & Y.Oshima, Species of the toxic dinoflagellate genus *Alexandrium* in southeastern Australian waters, *Bot. Mar.* 34: 575–587 (1991)

G.M.Hallegraeff, D.A.Steffensen & R.Wetherbee, Three estuarine Australian dinoflagellates that can produce paralytic shellfish toxins, *J. Plankton Res.* 10: 531–541 (1998)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)

P.Adjani, T.Ingleton, T.Pritchard & L.Armand, Microalgal blooms in the coastal waters of New South Wales, Australia, *Proc. Linn. Soc. New South Wales* 133: 15–31 (2011)

H.Farrell, S.Brett, P.Ajani & S.Murray, Distribution of the genus *Alexandrium* (Halim) and paralytic shellfish toxins along the coastline of New South Wales, Australia, *Mar. Poll. Bull.* 72: 133–145 (2013)