

Alexandrium margalefii Balech

Trans. Amer. Microscop. Soc. 113: 220, figs 12–15 (1994)

Alexandrium margalefi Balech, in G.M.Hallegraeff *et al.*, *Bot. Mar.* 34: 580 (1991), *nom. inval.*

Type locality: Spain

Australian distribution: N.S.W. (Central & South Coasts). Common in southern Australian coastal waters during warmer months; River Derwent and Georges Bay (Tas.), Port Phillip Bay (Vic.) and Port Lincoln (S.A.) [Hallegraeff *et al.*, 2010]

Extra-Australian distribution: Mexico, Spain, Denmark, New Zealand

References: G.M.Hallegraeff, C.J.[S.]Bolch, S.I.Blackburn & Y.Oshima, Species of the toxic dinoflagellate genus *Alexandrium* in southeastern Australian waters, *Bot. Mar.* 34: 575–587 (1991)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

H.Farrell, S.Brett, P.Ajani & S.Murray, Distribution of the genus *Alexandrium* (Halim) and paralytic shellfish toxins along the coastline of New South Wales, Australia, *Mar. Poll. Bull.* 72: 133–145 (2013)