

Ankistrodinium semilunatum (Herdman) Hoppenrath, Murray, Sparmann & Leander

J. Phycol. 48: 1145, figs 1–4 (2012)

Amphidinium semilunatum Herdman, *Trans. Proc. Liverpool Biol. Soc.* 38: 59, fig. 7 (1924)

Type locality: Great Britain

Australian distribution: Qld (Bowling Green Bay), N.S.W. (Botany Bay, Durras Lake, Narrabeen Lagoon, Sydney, Chowder Bay; in sand), W.A. (Shark Bay)

Extra-Australian distribution: U.S.A. (Alaska), Canada (British Columbia), British Isles, northern France, Wadden Sea (Denmark, Germany), Baltic Sea, Italy, Greece, Kuwait

References: J.Larsen & D.J.Patterson, Some flagellates (Protista) from tropical marine sediments, *J. Nat. Hist. (London)* 24: 801–937 (1990)

S.Murray & D.J.Patterson, The benthic dinoflagellate genus *Amphidinium* in south-eastern Australian waters, including 3 new species, *Eur. J. Phycol.* 37: 279–298 (2002)

S.Al-Qassab, W.J.Lee, S.Murray, A.G.B.Simpson & D.J.Patterson, Flagellates from stromatolites and surrounding sediments in Shark Bay, Western Australia, *Acta Protozool.* 41: 91–144 (2002)

S.Murray, Benthic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 213–259 (2010)

M.Hoppenrath, S.Murray, S.F.Sparmann & B.S.Leander, Morphology and molecular phylogeny of *Ankistrodinium* gen. nov. (Dinophyceae), a new genus of marine sand-dwelling dinoflagellates formerly classified within *Amphidinium*, *J. Phycol.* 48: 1143–1152 (2012)