

Ceratium azoricum Cleve

Kongl. Svenska Vetensk. Acad. Handl. 34: 13, pl. 7, figs 6, 7 (1900)

Neoceratium azoricum (Cleve) F.Gómez, D.Moreira & López-García, *Protist* 161: 43 (2010), *nom. illeg.*

Type locality: Azores

Ceratium petersii Steem.-Niels., *Dana Rep.* 4: 20, fig. 44 (1934)

Type locality: South Pacific Ocean

Australian distribution: East coast: Qld (Moreton Bay) to N.S.W. (Eden), Tas. (SE coast), W.A. (off Albany, W coast)

Extra-Australian distribution: South Atlantic Ocean, Gulf of Mexico, Azores, Canary Islands, U.S.A. (California, NE coast), North Atlantic Ocean, France, British Isles, Mediterranean Sea, Arabian Sea, Indian Ocean, Bay of Bengal, Andaman Sea, South China Sea, East China Sea, North Pacific Ocean, Solomon Islands, New Zealand, South Pacific Ocean, eastern tropical Pacific Ocean, Peru-Chile Current

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)