

Ceratium cephalotum (Lemmerm.) Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl.: 10, pl. 1, fig. 10 (1911)

Ceratium gravidum var. *cephalotum* Lemmerm., *Ber. Deutsch Bot. Ges.* 18: 349, pl. 1, fig. 15 (1900)

Neoceratium cephalotum (Lemmerm.) F.Gómez, D.Moreira & López-García, *Protist* 161: 43 (2010), *nom. illeg.*

Type locality: Atlantic Ocean [14°S, 27°W]

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld (Moreton Bay), N.S.W. (Trial Bay)

Extra-Australian distribution: NE Brazil, Mexico (Gulf of California), Gulf of Mexico, tropical Atlantic Ocean, Canary Islands, Réunion, Sri Lanka, Sumatra, Japan, western Pacific Ocean, Solomon Islands, South Pacific Ocean, eastern Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)