

Ceratium concilians Jörg.

Rep. Danish Oceanogr. Exped. Mediterr. Adjacent Seas 2, J.1: 72, fig. 69 (1920)

Neoceratium concilians (Jörg.) F.Gómez, D.Moreira & López-García, *Protist* 161: 43 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (Turkey)

Australian distribution: Qld (Coral Sea, Great Barrier Reef), N.S.W. (Trial Bay to Eden, off Sydney), Bass Strait, Tas.

Extra-Australian distribution: Gulf of Mexico, Canary Islands, Bay of Biscay, British Isles, Spain, Mediterranean Sea, Adriatic Sea, Red Sea, Arabian Sea, Solomon Islands, New Zealand, North Pacific Ocean, Tasman Sea, New Zealand, Antarctic waters, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)