

Ceratium deflexum (Kofoid) Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl. 2(1): 64 (1911)

Ceratium macroceros subsp. *deflexum* Kofoid, *Univ. Calif. Publ. Zool.* 3: 304, pl. 24, figs 13–15 (1907);
Neoceratium deflexum (Kofoid) F.Gómez, D.Moreira & López-García, *Protist* 161: 43 (2010), *nom. illeg.*

Type locality: U.S.A. (California)

Australian distribution: Coral Sea, Qld (Great Barrier Reef), N.S.W. (N coast), East Australian Current

Extra-Australian distribution: Brazil, tropical Atlantic Ocean, Gulf of Mexico, North Sea, Mediterranean Sea, Red Sea, Arabian Sea, Indian Ocean, Indonesia, Gulf of Thailand, South China Sea, East China Sea, Yellow Sea, Japan, North Pacific Ocean, SW Pacific Ocean, Pacific coast of Mexico, U.S.A. (California), Ecuador

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)