

Ceratium euarcuatum Jörg.

Rep. Danish Oceanogr. Exped. Mediterr. Adjacent Seas 2, J.1: 56, fig. 54 (1920)

Ceratium arcuatum (Gourret) Pavill. (1905), nom. illeg., non *C. arcuatum* Cleve (1900); *Neoceratium euarcuatum* (Jörg.) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), nom. illeg.

Type locality: Mediterranean Sea

Australian distribution: Qld (Great Barrier Reef), N.S.W., East Australian Current

Extra-Australian distribution: SW Africa, Brazil, Gulf of Mexico, U.S.A. (NE coast), tropical Atlantic Ocean, Mediterranean Sea, Red Sea, Arabian Sea, Indian Ocean, Indonesia, South China Sea, East China Sea, Solomon Islands, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)