

***Ceratium falcatifforme* Jörg.**

Rep. Danish Oceanogr. Exped. Mediterr. Adjacent Seas 2, J.1: 40, fig. 29 (1920)

Neoceratium falcatifforme (Jörg.) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (Spain)

Australian distribution: N.S.W., Bass Strait

Extra-Australian distribution: Brazil, Gulf of Mexico, Atlantic Ocean, Mediterranean Sea, Arabian Sea, tropical Pacific Ocean, Coral Sea, Tasman Sea, SW Pacific Ocean, New Zealand, Pacific coast of Mexico, SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)