

Ceratium gibberum Gourret

Ann. Mus. Hist. Nat. Marseille (Zool.) 1(8): 43, pl. 2, figs 33–35 (1883)

Neoceratium gibberum (Gourret) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (France)

?*Ceratium gibberum* f. *aequale* Jörg., *Rep. Danish Oceanogr. Exped. Mediterr. Adjacent Seas* 2, J.1: 70, fig. 68 (1920)

Type locality: Mediterranean Sea (Greece)

Australian distribution: Gulf of Carpentaria/North-west Shelf, E coast, Bass Strait

Extra-Australian distribution: Gulf of Mexico, Caribbean Sea, U.S.A. (Washington), NW Atlantic Ocean, Canary Islands, Iceland, British Isles, Bay of Biscay, North Sea, Mediterranean Sea (France, Spain, Italy, Greece), Adriatic Sea, Black Sea, Red Sea, Gulf of Guinea, East Africa, Arabian Sea, Indian Ocean, Bay of Bengal, Indonesia, Gulf of Thailand, Papua New Guinea, Solomon Islands, Japan, Coral Sea, New Caledonia, Tasman Sea, New Zealand, South Pacific Ocean, North Pacific Ocean, Pacific coast of South America

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)