

**Ceratium hexacanthum** Gourret

*Ann. Mus. Hist. Nat. Marseille (Zool.)* 1(8): 36, pl. 3, fig. 49, 49A (1883)

*Neoceratium hexacanthum* (Gourret) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), *nom. illeg*

Type locality: Mediterranean Sea (France)

Australian distribution: Qld (Coral Sea, Great Barrier Reef), N.S.W., Lord Howe Island, East Australian Current, Bass Strait, Tas., S.A., W.A.

Extra-Australian distribution: Brazil, Gulf of Mexico, U.S.A. (Florida), Atlantic Ocean, Canary Islands, British Isles, North Sea, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, Indian Ocean, Indonesia, East China Sea, Japan, Solomon Islands, Tasman Sea, New Zealand, Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)