

Ceratium incisum (G.Karst.) Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl.: 40, figs 82, 83 (1911)

Ceratium furca var. *incisum* G.Karst., *Wiss. Ergeb. Deutschen Tiefsee-Exped. "Valdivia"* 2(2): 149, pl. 23, fig. 6 (1906)

Type locality: Atlantic Ocean (01°56'S, 07°48'E)

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, N.S.W. (Port Hacking, Eden), East Australian Current

Extra-Australian distribution: Brazil, Caribbean Sea, Gulf of Mexico, Atlantic Ocean, Canary Islands, West Africa, Réunion, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, Indian Ocean, Andaman Sea, South China Sea, East China Sea, Japan, North Pacific Ocean, Solomon Islands, SW Pacific Ocean, eastern tropical Pacific Ocean, Peru

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)