

Ceratium inflatum (Kofoid) Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl. 2(1): 25 (1911)

Ceratium pennatum f. *inflatum* Kofoid, *Bull. Mus. Compar. Zool. Harvard Coll.* 50: 172, pl. 2, fig. 13 (1907);
Neoceratium inflatum (Kofoid) F.Gómez, D.Moreira & López-García, *Protist* 161: 44 (2010), *nom. illeg.*

Type locality: eastern tropical Pacific Ocean

Australian distribution: Qld (Great Barrier Reef), N.S.W. (near Sydney)

Extra-Australian distribution: Brazil, South Atlantic Ocean, South Africa, Caribbean Sea, Gulf of Mexico, U.S.A. (Alabama, Florida), North Atlantic Ocean, Canary Islands, Azores, British Isles, Mediterranean Sea, Black Sea, Red Sea, Arabian Sea, Indian Ocean, Bay of Bengal, Indonesia, Gulf of Thailand, South China Sea, East China Sea, Japan, North Pacific Ocean, New Zealand, eastern tropical Pacific Ocean, Peru, SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)