

**Ceratium lunula** (Schimp. ex G.Karst.) Jörg.

*Internat. Rev. Ges. Hydrobiol. Hydrogr.* 4, Biol. Suppl.: figs 112–115 (1911)

*Ceratium tripos* [var.] *lunula* Schimp. ex G.Karst., *Wiss. Ergeb. Deutschen Tiefsee-Exped. "Valdivia"* 2(2): 102, pl. 20, fig. 12 (1906); *Neoceratium lunula* (Schimp. ex G.Karst.) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: South Africa

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld (Great Barrier Reef, SE coast), N.S.W., east Australian Current, Vic. (S of Port Fairy), Bass Strait, Great Australian Bight, W.A.

Extra-Australian distribution: Brazil, South Africa, Gulf of Mexico, U.S.A. (NE coast), Canary Islands, Spain, British Isles, red Sea, Arabian Sea, Indian Ocean, South China Sea, East China Sea, Yellow Sea, Papua New Guinea, Solomon Islands, Coral Sea, Tasman Sea, New Zealand, South Pacific Ocean, Peru

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)