

Ceratium paradoxides Cleve

Kongl. Svenska Vetensk. Acad. Handl. 34: 15, pl. 7, fig. 1 (1900)

Neoceratium paradoxides (Cleve) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: Atlantic Ocean (Newfoundland Banks, Azores, Canary Islands)

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Qld (Great Barrier Reef, Moreton Bay) to N.S.W. (Sydney), Bass Strait

Extra-Australian distribution: Brazil, South Atlantic Ocean, Gulf of Mexico, Canada (Newfoundland Banks), Azores, Canary Islands, Mediterranean Sea, Arabian Sea, East Africa, Indian Ocean, SW Pacific Ocean, North Pacific Ocean, Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)