

Ceratium platycorne Daday

Termeszettud. Fuz. 11: 78, 101, pl. 3, figs 1, 2 (1888), as *platicornis*

Neoceratium platycorne (Daday) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: Mediterranean Sea (Italy)

Ceratium compressum auct. non Gran: E.J.F.Wood, *Austral. J. Mar. Freshw. Res.* 5: 312 (1954)

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld (Great Barrier Reef), N.S.W., East Australian Current, Vic., Bass Strait, Tas. (off E coast)

Extra-Australian distribution: Brazil, Gulf of Mexico, Gulf Stream, North Atlantic Ocean, Canary Islands, British Isles, Mediterranean Sea, Indian Ocean, Indonesia, South China Sea, East China Sea, Japan, Pacific Ocean, Tasman Sea, New Zealand, U.S.A. (California)

References: W.J.Dakin & A.N.Colefax, The marine plankton of the coastal waters of New South Wales. I. The chief planktonic forms and their seasonal distribution, *Proc. Linn. Soc. New South Wales* 58: 186–222 (1933)

E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

J.M.Huisman, The genus *Ceratium* Schrank (Dinophyceae) in Bass Strait and adjoining waters, southern Australia, *Austral. Syst. Bot.* 2: 425–454 (1989)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)