

Ceratium porrectum G.Karst.

Wiss. Ergeb. Deutsch. Tiefsee-Exped. "Valdivia", 2(2): 309, pl. 51, fig. 6 (1907)

Neoceratium porrectum (G.Karst.) F.Gómez, D.Moreira & López-García, *Protist* 161: 45 (2010), *nom. illeg.*

Type locality: Seychelles

Australian distribution: Qld (Great Barrier Reef), N.S.W. (Port Hacking), East Australian Current, Tas. (Storm Bay, Port Arthur)

Extra-Australian distribution: Brazil, Portugal, Mediterranean Sea (France), Seychelles, Indian Ocean (S of India), Pacific coast of Mexico

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)