

Ceratium praelongum (Lemmerm.) Kofoid ex Jörg.

Internat. Rev. Ges. Hydrobiol. Hydrogr. 4, Biol. Suppl.: (1911)

Ceratium gravidum var. *praelongum* Lemmerm., *Abh. Naturwiss. Vereine Bremen* 16: 349, pl. 1, fig. 15 (1899)

Type locality: Atlantic Ocean (14°S, 27°W)

Extra-Australian distribution: Gulf of Mexico, Caribbean Sea, Atlantic Ocean, Canary Islands, Mediterranean Sea, Red Sea, Indian Ocean, Bay of Bengal, Peru

Australian distribution: Gulf of Carpentaria/North-west Shelf, Coral Sea, Qld (Great Barrier Reef), East Australian Current

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

S.W.Jeffrey & G.M.Hallegraeff, Phytoplankton pigments, species and light climate in a complex warm-core eddy of the East Australian Current, *Deep Sea Res., Part A* 34: 649–673 (1987)