

***Dinophysis acuminata*** Clap. & J.Lachm.

*Mém. Inst. Nat. Genève.* 6 (1858): 408, pl. 20, fig. 17 (1859)

*Type locality:* North Sea (Norway)

*Australian distribution:* N.S.W. (Port Hacking, Twofold Bay), Vic., Tas. (River Derwent), S.A., W.A. (Swan River)

*Extra-Australian distribution:* U.S.A. (Washington), Canada (E coast), Canary Islands, British Isles, North Sea, Norwegian Sea, Baltic Sea, Mediterranean Sea (Turkey), Adriatic Sea, Japan, New Zealand

*Note:* A seasonal, bloom-forming species [Hallegraeff *et al.*, 2010]

*References:* E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

P.Adjani, T.Ingleton, T.Pritchard & L.Armand, Microalgal blooms in the coastal waters of New South Wales, Australia, *Proc. Linn. Soc. New South Wales* 133: 15–31 (2011)