

Dinophysis fortii Pavill.

Bull. Soc. Bot. France 70: 881 (1923)

Type locality: Mediterranean Sea (France)

Australian distribution: N.S.W. (Port Stephens to Jervis Bay), East Australian Current, Vic. (Port Fairy), Tas. (Storm Bay, River Derwent)

Extra-Australian distribution: Brazil, U.S.A. (Washington), British Isles, Mediterranean Sea (France), Adriatic Sea, Black Sea, Japan, New Zealand, Southern Ocean

Note: A potentially harmful species

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)