

Dinophysis tripos Gourett

Ann. Mus. Hist. Nat. Marseille (Zool.) 1(8): pl. 3, fig. 53 (1883)

Type locality: Mediterranean Sea (France)

Australian distribution: East Australian Current, N.S.W. (Sydney and offshore), Vic.

Extra-Australian distribution: Brazil, Gulf Stream, North Atlantic Ocean, British Isles, Norwegian Sea, Greenland, Mediterranean Sea, SE Asia, Japan, New Zealand

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)