

**Diplopsalopsis bomba** (F.Stein ex Jörg.) J.D.Dodge & S.Toriumi

*Bot. Mar.* 36: 145 (1993), *nom. inval.*

*Diplopelta bomba* F.Stein ex Jörg. (1912)

*Type locality:* ?

*Peridiniopsis asymmetrica* Mangin, *Compt. Rend. Hebd. Séances Acad. Sci.*, sér. D, 153:31, fig. 2:1, 2 (1911);  
*Dissodium asymmetricum* (Mangin) A.R.Loebl., *Proc. N. Amer. Paleont. Conv. Chicago*, 1969 905 (1970)

*Type locality:* France (Atlantic coast)

*Australian distribution:* Gulf of Carpentaria/North-west Shelf, N.S.W. (off Sydney)

*Extra-Australian distribution:* Brazil, Gulf of Mexico, U.S.A. (Virginia), Canada (Gulf of St. Lawrence), British Isles, North Sea, Baltic Sea, Arctic Ocean, France, Mediterranean Sea, Black Sea, Bay of Bengal, South China Sea, Japan, Peru

*References:* G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)