

Goniodoma sphaericum G.Murray & Whitting

Trans. Linn. Soc. London, Ser. Bot. 5: 335, pl. 27, fig. 3 ('1899') [1900]

Triadinium sphaericum (G.Murray & Whitting) J.D.Dodge, *Brit. Phycol. J.* 16: 279 (1981)

Type locality: Atlantic Ocean

Australian distribution: Gulf of Carpentaria/North-west Shelf, East Australian Current , W of Bass Strait

Extra-Australian distribution: Gulf of Mexico, Atlantic Ocean, Canary Islands, British Isles, Spain, Mediterranean Sea, Indian Ocean, Singapore, Mexico (Baja California)

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

S.W.Jeffrey & G.M.Hallegraeff, Studies of phytoplankton species and photosynthetic pigments in a warm core eddy of the East Australian Current. I. Summer populations, *Mar. Ecol. Prog. Ser.* 3: 285–294 (1980)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)